

On a clean determination of ϕ_2

A. I. SANDA

Department of Physics, Nagoya University
Nagoya, Japan

Abstract

We point out that time dependent study of four decay modes $B^0(t) \rightarrow \overline{D}K_S$, $B^0(t) \rightarrow DK_S$, $\overline{B}^0(t) \rightarrow \overline{D}K_S$, and $\overline{B}^0(t) \rightarrow DK_S$ allows us to measure $\sin(\phi_1 - \phi_2)$ - contrary to previously obtained result that it measures $\sin(2\phi_1 + \phi_3)$. The time dependent study of $B_s^0(t) \rightarrow D_s^\pm K^\mp$, and the study of $B^- \rightarrow K^-(D, \overline{D}) \rightarrow K^- f$ decay where $D \rightarrow f$ is a doubly Cabibbo suppressed decay yield information on ϕ_3 , as previously expected.

1. Introduction

Large **CP** violation in $B^0 \rightarrow \psi K_S$ has been observed by Babar and Belle¹. Clean measurement of ϕ_1 has been made. This is the first step toward a serious test of the standard model of **CP** violation. The statistical and systematic errors are large but this will improve in due time. New CDF measurements are around the corner. LHCb and BTeV is also coming up. Exciting program for **CP** violation study is on its way. What is next? We would like to test the unitarity relation $\phi_1 + \phi_2 + \phi_3 = \pi$. How do we measure ϕ_2 ? For many years, **CP** asymmetry in $B^0 \rightarrow \pi\pi$ was thought to be the way to measure ϕ_2 . The existence of penguin amplitudes which prohibit a theoretically clean measurement was overcome by the isospin analysis. But, recent indication² that $Br(B^0 \rightarrow \pi^0\pi^0) \sim 2 \times 10^{-7}$, and the background associated with π^0 detection makes it very difficult to extract ϕ_2 from the isospin analysis.

A Dalitz plot analysis of $B \rightarrow \rho\pi \rightarrow 3\pi$ has been proposed³. While we expect $B \rightarrow 3\pi$ to be dominated by $B \rightarrow \rho\pi$ which is a pure **CP** eigenstate, there is some opposite **CP** admixture under the ρ band in the Dalitz plot. Further study of the Dalitz plot is necessary to see if this

technique can be used to extract ϕ_2 without theoretical ambiguity.

2. Determination of ϕ_2 from $B^0, \bar{B}^0 \rightarrow DK_S$ Decay

Here we ask if there is a practical way to determine ϕ_2 in a theoretically clean way. We point out that $\sin(\phi_1 - \phi_2)$ can be determined without theoretical ambiguity by studying two time dependent asymmetries:

$$\begin{aligned} a(t) &= \frac{\Gamma(\bar{B}^0(t) \rightarrow DK_S) - \Gamma(B^0(t) \rightarrow DK_S)}{\Gamma(\bar{B}^0(t) \rightarrow DK_S) + \Gamma(B^0(t) \rightarrow DK_S)} \\ \bar{a}(t) &= \frac{\Gamma(\bar{B}^0(t) \rightarrow \bar{D}K_S) - \Gamma(B^0(t) \rightarrow \bar{D}K_S)}{\Gamma(\bar{B}^0(t) \rightarrow \bar{D}K_S) + \Gamma(B^0(t) \rightarrow \bar{D}K_S)}. \end{aligned} \quad (1)$$

Here, in addition to $B^0 \rightarrow DK_S$, we can also use $B^0 \rightarrow DK^{0*} \rightarrow DK_S \pi^0$, and $B^0 \rightarrow D^{0*} K_S$.

In the literature⁴, it is stated that this asymmetry measures $\sin(2\phi_1 + \phi_3)$ which of course is identical to $-\sin(\phi_1 - \phi_2)$ if we assume unitarity, $\phi_1 + \phi_2 + \phi_3 = \pi$. Another often made statement is that $\arg(\mathbf{V}_{ub}^* \mathbf{V}_{ud} \mathbf{V}_{tb} \mathbf{V}_{td}^*) = \phi_1 + \phi_3$. The correct statement is that it is related to ϕ_2 as given in Eq. (3). One should remember not to use equalities which depend on a particular phase convention, for example, $\arg(\mathbf{V}_{ub}^* \mathbf{V}_{ud}) = \phi_3$ or $\arg(\mathbf{V}_{tb} \mathbf{V}_{td}^*) = \phi_1$. In the first round of theoretical analysis, we were mainly interested in identifying all possible **CP** asymmetries, and we did not exercise sufficient care in deriving the expression for **CP** asymmetry independent of the unitarity. The unitarity constraint, which we want to test, has crept into derivations.

To avoid using unitarity relation which we want to test, it is useful to express all physical observables in a rephasing invariant way, and write the asymmetries in terms of ϕ_1, ϕ_2, ϕ_3 using the definitions:

$$\phi_1 = \pi - \arg\left(\frac{-\mathbf{V}_{tb}^* \mathbf{V}_{td}}{-\mathbf{V}_{cb}^* \mathbf{V}_{cd}}\right), \quad (2)$$

$$\phi_2 = \arg\left(\frac{\mathbf{V}_{tb}^* \mathbf{V}_{td}}{-\mathbf{V}_{ub}^* \mathbf{V}_{ud}}\right), \quad (3)$$

$$\phi_3 = \arg\left(\frac{\mathbf{V}_{ub}^* \mathbf{V}_{ud}}{-\mathbf{V}_{cb}^* \mathbf{V}_{cd}}\right). \quad (4)$$




Figure 1: The unitarity triangle in the complex plane.

The **CP** asymmetry for $\bar{B}^0 \rightarrow \psi K_S$ is theoretically clean at the level where we neglect $\mathcal{O}(\sin^2 \theta_c)$ terms. Here θ_c is the Cabibbo angle. This comes from the fact that, for $B^0 \rightarrow \psi K_S$ decay, the tree graph, which is proportional to $\mathbf{V}_{cb} \mathbf{V}_{cs}^*$, and the penguin graph, which is proportional to $\mathbf{V}_{tb} \mathbf{V}_{ts}^*$, have the same phase if we neglect terms proportional to $\mathbf{V}_{ub} \mathbf{V}_{us}^*$. The latter neglected term is down by $\mathcal{O}(\sin^2 \theta_c)$ compared to the former.

The 2×2 submatrices of the KM matrix are known to be approximately unitary:

$$\begin{aligned}\mathbf{V}_{ud}^* \mathbf{V}_{us} &\cong -\mathbf{V}_{cd}^* \mathbf{V}_{cs} \\ \mathbf{V}_{ud} \mathbf{V}_{cd}^* &\cong -\mathbf{V}_{us} \mathbf{V}_{cs}^* \\ \mathbf{V}_{cs}^* \mathbf{V}_{cb} &\cong -\mathbf{V}_{ts}^* \mathbf{V}_{tb}.\end{aligned}\tag{5}$$

These relations have been verified experimentally to within 6%, .1%, 3%, respectively, relative to terms that are kept⁵. So, since we are making statements that are accurate to order $\sin^2 \theta_c$, we are free to use these relations to test the unitarity relation:

$$\mathbf{V}_{td} \mathbf{V}_{ud}^* + \mathbf{V}_{ts} \mathbf{V}_{us}^* + \mathbf{V}_{tb} \mathbf{V}_{ub}^* = 0\tag{6}$$

Now, let us discuss $B^0(t) \rightarrow \overline{D}K_S$, $B^0(t) \rightarrow DK_S$, $\overline{B}^0(t) \rightarrow \overline{D}K_S$, and $\overline{B}^0(t) \rightarrow DK_S$ decays. Defining $|K_S\rangle$ state as

$$|K_S\rangle = p_K |K^0\rangle + q_K |\overline{K}^0\rangle\tag{7}$$

and using **CP** symmetry of strong interaction, we obtain

$$\begin{aligned}A(\overline{B} \rightarrow \overline{D}K_S) &= e^{i\delta_-} \mathbf{V}_{cb} \mathbf{V}_{us}^* \mathcal{A}_- q_K^* \\ A(\overline{B} \rightarrow DK_S) &= e^{i\delta_+} \mathbf{V}_{ub} \mathbf{V}_{cs}^* \mathcal{A}_+ q_K^* \\ A(B \rightarrow DK_S) &= e^{i\delta_-} \mathbf{V}_{cb}^* \mathbf{V}_{us} \mathcal{A}_- p_K^* \\ A(B \rightarrow \overline{D}K_S) &= e^{i\delta_+} \mathbf{V}_{ub}^* \mathbf{V}_{cs} \mathcal{A}_+ p_K^*,\end{aligned}\tag{8}$$

where $\delta_{\pm}$ are strong phases, $\mathcal{A}_{\pm}$ are the magnitudes of the matrix elements.

Note that $|A(\overline{B} \rightarrow DK_S)| = |A(B \rightarrow DK_S)|$ obviously does *not* hold as an identity. Setting $\Delta\Gamma_B = 0$, for simplicity, we find

$$\begin{aligned}\Gamma(B(t) \rightarrow DK_S) &\propto |A(DK_S)|^2 \left[1 + |\overline{\rho}(DK_S)|^2 \right. \\ &\quad \left. + (1 - |\overline{\rho}(DK_S)|^2) \cos \Delta M_B t - 2 \text{Im} \left(\frac{q_B}{p_B} \overline{\rho}(DK_S) \right) \sin \Delta M_B t \right] \\ \Gamma(\overline{B}(t) \rightarrow \overline{D}K_S) &\propto |\overline{A}(\overline{D}K_S)|^2 \left[1 + |\rho(\overline{D}K_S)|^2 \right. \\ &\quad \left. + (1 - |\rho(\overline{D}K_S)|^2) \cos \Delta M_B t - 2 \text{Im} \left(\frac{p_B}{q_B} \rho(\overline{D}K_S) \right) \sin \Delta M_B t \right] \\ \Gamma(B(t) \rightarrow \overline{D}K_S) &\propto |A(\overline{D}K_S)|^2 \left[1 + |\overline{\rho}(\overline{D}K_S)|^2 \right. \\ &\quad \left. + (1 - |\overline{\rho}(\overline{D}K_S)|^2) \cos \Delta M_B t - 2 \text{Im} \left(\frac{q_B}{p_B} \overline{\rho}(\overline{D}K_S) \right) \sin \Delta M_B t \right] \\ \Gamma(\overline{B}(t) \rightarrow DK_S) &\propto |\overline{A}(DK_S)|^2 \left[1 + |\rho(DK_S)|^2 \right. \\ &\quad \left. + (1 - |\rho(DK_S)|^2) \cos \Delta M_B t - 2 \text{Im} \left(\frac{p_B}{q_B} \rho(DK_S) \right) \sin \Delta M_B t \right]\end{aligned}\tag{9}$$

Using

$$\frac{q_B q_K^*}{p_B p_K^*} = \frac{\mathbf{V}_{td} \mathbf{V}_{tb}^* \mathbf{V}_{cs} \mathbf{V}_{cd}^*}{\mathbf{V}_{td}^* \mathbf{V}_{tb} \mathbf{V}_{cs}^* \mathbf{V}_{cd}}.\tag{10}$$

we obtain:

$$\begin{aligned}\frac{q_B}{p_B}\bar{\rho}(\overline{D}K_S) &= e^{i(\delta_- - \delta_+)} \frac{\mathbf{V}_{cb}\mathbf{V}_{us}^*}{\mathbf{V}_{ub}^*\mathbf{V}_{cs}} \frac{\mathbf{V}_{td}\mathbf{V}_{tb}^*}{\mathbf{V}_{td}^*\mathbf{V}_{tb}} \frac{\mathbf{V}_{cs}\mathbf{V}_{cd}^*}{\mathbf{V}_{cs}^*\mathbf{V}_{cd}} \frac{\mathcal{A}_-}{\mathcal{A}_+} \\ \frac{q_B}{p_B}\bar{\rho}(DK_S) &= e^{-i(\delta_- - \delta_+)} \frac{\mathbf{V}_{ub}\mathbf{V}_{cs}^*}{\mathbf{V}_{cb}^*\mathbf{V}_{us}} \frac{\mathbf{V}_{td}\mathbf{V}_{tb}^*}{\mathbf{V}_{td}^*\mathbf{V}_{tb}} \frac{\mathbf{V}_{cs}\mathbf{V}_{cd}^*}{\mathbf{V}_{cs}^*\mathbf{V}_{cd}} \frac{\mathcal{A}_+}{\mathcal{A}_-}\end{aligned}\quad (11)$$

Note that this is rephasing invariant. Using Eq. (5), we obtain:

$$\frac{\mathbf{V}_{cb}\mathbf{V}_{us}^*}{\mathbf{V}_{ub}^*\mathbf{V}_{cs}} \frac{\mathbf{V}_{td}\mathbf{V}_{tb}^*}{\mathbf{V}_{td}^*\mathbf{V}_{tb}} \frac{\mathbf{V}_{cs}\mathbf{V}_{cd}^*}{\mathbf{V}_{cs}^*\mathbf{V}_{cd}} = -\frac{\mathbf{V}_{cb}\mathbf{V}_{cd}^*}{\mathbf{V}_{ub}^*\mathbf{V}_{ud}} \frac{\mathbf{V}_{td}\mathbf{V}_{tb}^*}{\mathbf{V}_{td}^*\mathbf{V}_{tb}} \quad (12)$$

and using Eq. (4):

$$\begin{aligned}\frac{q_B}{p_B}\bar{\rho}(\overline{D}K_S) &= -e^{i(\delta_- - \delta_+)} \frac{\mathcal{A}_-}{\mathcal{A}_+} \frac{1}{\mathcal{R}} e^{i(\phi_2 - \phi_1)} \\ \frac{q_B}{p_B}\bar{\rho}(DK_S) &= -e^{-i(\delta_- - \delta_+)} \frac{\mathcal{A}_+}{\mathcal{A}_-} \mathcal{R} e^{i(\phi_2 - \phi_1)}\end{aligned}\quad (13)$$

where

$$\mathcal{R} = \left| \frac{\mathbf{V}_{ub}\mathbf{V}_{ud}^*}{\mathbf{V}_{cb}^*\mathbf{V}_{cd}} \right| \quad (14)$$

Measurement of the coefficients of $\cos \Delta M_B t$, we can determine $\frac{\mathcal{A}_+}{\mathcal{A}_-} \mathcal{R}$. The coefficient of $\sin \Delta M_B t$ gives $\sin(\phi_1 - \phi_2)$.

3. Determination of ϕ_3 with $B^- \rightarrow K^- D$ decays

It is useful to reanalyze the other well known methods to determine ϕ_3 . Measurements of $Br(B^- \rightarrow K^-(D^0, \overline{D}^0) \rightarrow K^- f)$, the doubly Cabibbo suppressed decay $Br(D^0 \rightarrow f)$, $Br(\overline{D}^0 \rightarrow f)$, $Br(B^- \rightarrow K^- D^0)$, and $Br(B^- \rightarrow K^- \overline{D}^0)$ allow us to determine⁷ ϕ_3 . Consider

$$B^- \rightarrow K^-(D^0, \overline{D}^0) \rightarrow K^- f \quad (15)$$

Write

$$\begin{aligned}\mathcal{A} &= A(B^- \rightarrow D^0 K^-) A(D^0 \rightarrow f) \\ \overline{\mathcal{A}} &= A(B^- \rightarrow \overline{D}^0 K^-) A(\overline{D}^0 \rightarrow f)\end{aligned}\quad (16)$$

where these amplitudes represent strong interaction part and the KM factors are taken out.

$$A(B^- \rightarrow K^-(D^0, \overline{D}^0) \rightarrow f K^-) = [|\mathcal{A}|e^{i\delta}\mathbf{V}_{cb}\mathbf{V}_{us}^*\mathbf{V}_{us}\mathbf{V}_{cd}^* + |\overline{\mathcal{A}}|e^{i\bar{\delta}}\mathbf{V}_{ub}\mathbf{V}_{cs}^*\mathbf{V}_{ud}^*\mathbf{V}_{cs}] \quad (17)$$

Here δ and $\bar{\delta}$ are strong interaction phases for decays involving D and $\overline{D}$ intermediate particles, respectively. It should be noted that by choosing the doubly Cabibbo suppressed decay of D meson, and that $B^- \rightarrow K^- \overline{D}$ is color suppressed, the two terms on the right hand side have approximately equal magnitude.




Figure 2: $A_1 = |A(B^- \rightarrow K^-(D^0, \bar{D}^0) \rightarrow K^- f)|$, $A_2 = |A(B^- \rightarrow D^0 K^-)A(D^0 \rightarrow f)|$ and $A_3 = |A(B^- \rightarrow \bar{D}^0 K^-)A(\bar{D}^0 \rightarrow f)|$ form a triangle. The angle at the top of the triangle $\phi = -\phi_3 + \bar{\delta} - \delta$. The analysis of the charge conjugate modes give $\bar{\phi} = -\phi_3 - \bar{\delta} + \delta$. Thus ϕ_3 can be determined.

The relative phase of these amplitudes is given by

$$\frac{A(B^- \rightarrow K^- D^0 \rightarrow K^- f)}{A(B^- \rightarrow K^- \bar{D}^0 \rightarrow K^- f)} = \frac{e^{i\delta} \mathbf{V}_{cb} \mathbf{V}_{us}^* \mathbf{V}_{cd}^* \mathbf{V}_{us}}{e^{i\bar{\delta}} \mathbf{V}_{ub} \mathbf{V}_{cs}^* \mathbf{V}_{ud}^* \mathbf{V}_{cs}} \frac{|\mathcal{A}|}{|\bar{\mathcal{A}}|} \quad (18)$$

Noting that

$$\begin{aligned} A_1 &= \sqrt{Br(B^- \rightarrow K^-(D^0, \bar{D}^0) \rightarrow K^- f)}, \\ A_2 &= \sqrt{Br(B^- \rightarrow D^0 K^-) Br(D^0 \rightarrow f)} \propto |\mathcal{A} \mathbf{V}_{cb} \mathbf{V}_{us}^* \mathbf{V}_{us} \mathbf{V}_{cd}^*| \\ A_3 &= \sqrt{Br(B^- \rightarrow \bar{D}^0 K^-) Br(\bar{D}^0 \rightarrow f)} \propto |\bar{\mathcal{A}} \mathbf{V}_{ub} \mathbf{V}_{cs}^* \mathbf{V}_{ud}^* \mathbf{V}_{cs}|, \end{aligned} \quad (19)$$

we can draw a triangle whose base is A_1 and the two sides are given by A_2 and A_3 . Eq. (18) gives the angle at the top of the triangle:

$$\phi = \pi - \arg \left[\frac{e^{i\delta} \mathbf{V}_{cb} \mathbf{V}_{cd}^*}{e^{i\bar{\delta}} \mathbf{V}_{ub} \mathbf{V}_{ud}^*} \right] = -\phi_3 + \bar{\delta} - \delta \quad (20)$$

The same analysis for B^+ decay leads to the determination of $\phi_3 + \bar{\delta} - \delta$ so both ϕ_3 and the relative strong interaction phase $\delta - \bar{\delta}$ can be determined.

4. Determination of ϕ_3 with $B_s^0 \rightarrow D_s^\pm K^\mp$ decays

Both $B_s \rightarrow D_s^\pm K^\mp$ and $\bar{B}_s^0 \rightarrow D_s^\pm K^\mp$ exist and there will be **CP** asymmetry which is expected to lead to information on ϕ_3 . We start by denoting relevant amplitudes:

$$\begin{aligned} A(\bar{B}_s \rightarrow D_s^- K^+) &= e^{i\delta_-} \mathbf{V}_{ub} \mathbf{V}_{cs}^* \mathcal{A}_- \\ A(\bar{B}_s \rightarrow D_s^+ K^-) &= e^{i\delta_+} \mathbf{V}_{cb} \mathbf{V}_{us}^* \mathcal{A}_+ \\ A(B_s \rightarrow D_s^- K^+) &= e^{i\delta_+} \mathbf{V}_{cb}^* \mathbf{V}_{us} \mathcal{A}_+ \\ A(B_s \rightarrow D_s^+ K^-) &= e^{i\delta_-} \mathbf{V}_{ub}^* \mathbf{V}_{cs} \mathcal{A}_-, \end{aligned} \quad (21)$$

where $\delta_\pm$ are strong phases, $\mathcal{A}_\pm$ are the magnitudes of the matrix elements.

$$\frac{q_{B_s} A(\bar{B}_s \rightarrow D_s^+ K^-)}{p_{B_s} A(B_s \rightarrow D_s^+ K^-)} = \frac{e^{i\delta_+} \mathbf{V}_{tb}^* \mathbf{V}_{ts}}{e^{i\delta_-} \mathbf{V}_{tb} \mathbf{V}_{ts}^*} \frac{\mathbf{V}_{us}^* \mathbf{V}_{cb}}{\mathbf{V}_{cs} \mathbf{V}_{ub}^*} \frac{\mathcal{A}_+}{\mathcal{A}_-} \quad (22)$$

$$\begin{aligned}
\frac{\mathbf{V}_{tb}^* \mathbf{V}_{ts}}{\mathbf{V}_{tb} \mathbf{V}_{ts}^*} \frac{\mathbf{V}_{us}^* \mathbf{V}_{cb}}{\mathbf{V}_{cs} \mathbf{V}_{ub}^*} &\cong \frac{\mathbf{V}_{cb}^* \mathbf{V}_{cs}}{\mathbf{V}_{cb} \mathbf{V}_{cs}^*} \frac{\mathbf{V}_{us}^* \mathbf{V}_{cb}}{\mathbf{V}_{cs} \mathbf{V}_{ub}^*} \\
&= \frac{\mathbf{V}_{cb}^* \mathbf{V}_{us}^* \mathbf{V}_{cs} \mathbf{V}_{ud}}{\mathbf{V}_{cs}^* \mathbf{V}_{ub}^* \mathbf{V}_{cs} \mathbf{V}_{ud}} \\
&\cong -\frac{\mathbf{V}_{cb}^* \mathbf{V}_{cd} \mathbf{V}_{ud}^* \mathbf{V}_{ud}}{\mathbf{V}_{cs}^* \mathbf{V}_{cs} \mathbf{V}_{ub}^* \mathbf{V}_{ud}} \\
&\cong \frac{1}{\mathcal{R}} e^{-i\phi_3}
\end{aligned} \tag{23}$$

$$\frac{q_{B_s}}{p_{B_s}} \frac{A(\overline{B}_s \rightarrow D_s^+ K^-)}{A(B_s \rightarrow D_s^+ K^-)} = \frac{1}{\mathcal{R}} \frac{\mathcal{A}_+}{\mathcal{A}_-} e^{i(\delta_+ - \delta_-) - i\phi_3} \tag{24}$$

Using Eq. (5), the right hand side of Eq. (25) can be written as Using the time dependence similar to Eq. (9), we can obtain both the magnitude and the phase of $\frac{q_{B_s}}{p_{B_s}} \frac{A(\overline{B}_s \rightarrow D_s^+ K^-)}{A(B_s \rightarrow D_s^+ K^-)}$. Similarly we can obtain the magnitude and the phase of

$$\frac{q_{B_s}}{p_{B_s}} \frac{A(\overline{B}_s \rightarrow D_s^- K^+)}{A(B_s \rightarrow D_s^- K^+)} = \frac{e^{i\delta_-} \mathbf{V}_{tb}^* \mathbf{V}_{ts} \mathbf{V}_{ub} \mathbf{V}_{cs}^* \mathcal{A}_-}{e^{i\delta_+} \mathbf{V}_{tb} \mathbf{V}_{ts}^* \mathbf{V}_{cb}^* \mathbf{V}_{us} \mathcal{A}_+} \tag{25}$$

$$= \mathcal{R} \frac{\mathcal{A}_-}{\mathcal{A}_+} e^{i(\delta_- - \delta_+) - i\phi_3}. \tag{26}$$

Thus ϕ_3 can be determined.

5. Summary

The time has come to perform a serious test of the KM ansatz for **CP** violation. To see if the unitarity triangle is actually a triangle is the next step. For this purpose we should be careful in deriving the expression for asymmetry in terms of three angles of the unitarity triangle. A useful rule is to always make sure that the asymmetry is indeopendent of phase conventions. We have reanalyzed processes known to yeild information on ϕ_3 . Asymmetries for B^0 , $\overline{B}^0 \rightarrow DK_S$ decays are shown to yield information on $\phi_2 - \phi_1$, and not on $2\phi_1 + \phi_3$ as previously stated. The other modes analyzed yield information on ϕ_3 as expected.

Acknowledgments

The author's research is supported by Grant-in Aid for Special Project Research (Physics of CP violation). I thank T. Kurimoto, Y. Okada, K. Ukai, and M. Yamauchi for useful discussions.

References

1. SLAC-PUB-8904 hep-ex/0107013; KEK Preprint 2001-50, Belle preprint 2001-10 , hep-ex/0107061
2. C. D. Lü, K. Ukai, and M. Z. Yang, *Phys. Rev.* **DD63** 074009 (2001).
3. H.R. Quinn and A.E. Snyder, *Phys. Rev.* **D48** 2139 (1993)

4. I. I. Bigi and A. I. Sanda *CP Violation* Cambridge, Cambridge University Press (1999).
5. F. J. Gilman, K. Kleinknecht, and B. Renk, Review of the CKM Quark Mixing Matrix in the 2000 Review of Particle Physics of D. E. Groom *et al.*, *Eur. J. Phys.* **C15**, 110-114 (2000).
6. D. Atwood, I. Dunietz and A. Soni, *Phys. Rev. Lett.* **78** 3257 (1997).